

Facial Fracture Management: Operative Trends

Ramon Garza, III MD

Lehigh Valley Health Network, Ramon.Garzaiii@lvhn.org

Joshua M. Adkinson MD

Lehigh Valley Health Network

Jarom N. Gilstrap MD

Lehigh Valley Health Network, Jarom_N.Gilstrap@lvhn.org

Sherrine Eid MPH

Lehigh Valley Health Network, Sherrine.Eid@lvhn.org

Robert X. Murphy Jr, MD, MS

Lehigh Valley Health Network, Robert.Murphy@lvhn.org

Follow this and additional works at: <https://scholarlyworks.lvhn.org/surgery>



Part of the [Other Medical Specialties Commons](#), and the [Surgery Commons](#)

Let us know how access to this document benefits you

Published In/Presented At

Garza, R., Adkinson, J., Gilstrap, J., Eid, S., & Murphy, R. (October 20-23, 2011). *Facial Fracture Management: Operative Trends*. Northeastern Society of Plastic Surgeons Annual Meeting, Amelia Island, Florida.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Facial Fracture Management: Operative Trends


Ramon Garza, III, M.D., Joshua M. Adkinson, M.D., Jarom N. Gilstrap, M.D., Sherrine Eid, M.P.H., Robert X. Murphy, Jr., M.D., M.S.
Lehigh Valley Health Network, Allentown, PA

Background:

Controversy exists in the literature regarding management of facial fractures. Indications for operative intervention can be subjective and may not reflect the clinical severity of the fracture or account for patient demographics. The objective of this study is to examine management of facial fractures at our Level 1 Trauma Center.

Methods:

- IRB-approved review of the Lehigh Valley Health Network Trauma Registry from 2000-2010
- Data Points Included: Age, gender, Injury Severity Score (ISS), Glasgow Coma Score (GCS), Facial Fractures (orbit, nasal, maxillary/malar, mandible), and Surgical Repair
- Logistic regression analysis was performed using SPSS 15.0 (SPSS Inc, Chicago, IL).



Results:

- A total 44,900 trauma patients were evaluated from 2000-2010
- 3,114 patients sustained 4,283 facial fractures
- Nasal Fractures: n = 1769 (41%)
- Malar/Maxillary Fractures: n = 1008 (24%)
- Orbital Fractures: n = 842 (20%)
- Mandible Fractures: n = 664 (15%)

Table 1. Likelihood of Operative Management of Facial Fractures (n=2938, p<0.001)

Independent Variables	p value	Adjusted OR (95% CI)
Age	<0.001	0.979 (0.975, 0.983)
Male	0.176	1.156 (0.937, 1.425)
Female		1.0 (reference)
ISS	0.394	0.996 (0.987, 1.005)
GCS	0.621	1.015 (0.982, 1.425)
Number of Fractures	<.001	2.604 (2.291, 2.960)

Figure 1. Facial Fracture Surgical Intervention (2000-2010)




Figure 2. Facial Fracture Inpatient Surgical Intervention (2000-2010)





Figure 3. Repair of Facial Fractures



References:

1. Evans BG, Evans GR. MOC-PSSM CME article: Zygomatic fractures. *Plast Reconstr Surg* 2008;121(1 Suppl):1-11.
2. Goldenberg, DV, Alonso, N, Ferreiro, M. Facial Trauma. In: Guyuron B, Eriksson, E, Persing, J (eds), (2009) *Plastic Surgery: Indications and Practice*. Saunders, New York, NY.
3. Iida S, Hassfeld S, Reuther T, et al. Maxillofacial fractures resulting from falls. *J Craniomaxillofac Surg* 2003;31(5):278-83.
4. Derdyn C, Persing JA, Broadus WC, et al. Craniofacial trauma: an assessment of risk related to timing of surgery. *Plast Reconstr Surg* 1990;86(2):238-45.
5. Shibuya TY, Karam AM, Doerr T, Stachler RJ, et al. Facial fracture repair in the traumatic brain injury patient. *J Oral Maxillofac Surg* 2007;65(9):1693-9.
6. Hollier Jr, L, Kelley, P. Soft Tissue and Skeletal Injuries of the Face. In: Aston, SJ, Beasley RW, Thorne CHM (eds), (2006) *Grabb and Smith's Plastic Surgery*. 6th Ed. Lippincott, Philadelphia, PA.
7. Sharabi SE, Koshy JC, Thornton JF, et al. Facial fractures (Review). *Plast Reconstr Surg* 2011;127(2):256-34e.
8. Thaller SR, Kawamoto HK. Care of maxillofacial injuries: Survey of plastic surgeons. *Plast Reconstr Surg* 1992;90(4):562-7.
9. Janus SC, MacLeod SP, Odland R. Analysis of results in early versus late midface fracture repair. *Otolaryngol Head Neck Surg* 2008;138(4):464-7.
10. Burnstine MA. Clinical recommendations for repair of orbital facial fractures. *Curr Opin Ophthalmol* 2003;14(5):236-40.
11. Covington DS, Wainwright DJ, Teichgraber JF, et al. Changing patterns in the epidemiology and treatment of zygoma fractures: 10-year review. *J Trauma* 1994;37(2):243-8.
12. Murphy RX Jr, Birmingham KL, Okunski WJ, et al. The influence of airbag and restraining devices on the patterns of facial trauma in motor vehicle collisions. *Plast Reconstr Surg* 2000;105(2):516-20.
13. Stacey DH, Doyle JF, Mount DL, et al. Management of mandible fractures [Review]. *Plast Reconstr Surg* 2006;117(3):486-60e.
14. Maloney PL, Welch TB, Doku HC. Early immobilization of mandibular fractures: A retrospective study. *J Oral Maxillofac Surg* 1991;49(7):698-702.
15. Gerbino G, Roccia F, De Giovanni PP, et al. Maxillofacial trauma in the elderly. *J Oral Maxillofac Surg* 1999;57(7):777-82.
16. Sanger C, Argenta LC, David LR. Cost-effective management of isolated facial fractures. *J Craniomaxillofac Surg* 2004;15(4):636-41.
17. Erdmann D, Price K, Reed S, et al. A financial analysis of operative facial fracture management. *Plast Reconstr Surg* 2008;121(4):1323-7.

Conclusion:

At our Level 1 Trauma Center, most patients with facial fractures are managed non-operatively. This trend seems to be increasing as the age of our patient population increases. Patients who have sustained multiple facial fractures, particularly involving the mandible, are more likely to undergo surgical repair. Additionally, there may be significant value to society in cost savings by non-operative management of patients with uncomplicated and clinically stable mid-facial fractures.