


# Infection

INFECTION CONTROL AND PREVENTION

# CONNECTION



VOLUME 6, ISSUE 1

JAN THRU MAR 2015

## IN THIS ISSUE:

- Collection Technique for Quantiferon-TB Gold Test .....2
- Disposable Wipes.....2
- Droplet Precautions - Personal Protective Equipment (PPE).....1
- Ebola.....1
- Fit Test Products.....2
- Linen Hampers Required for Contact Isolation.....2
- N-95 and PAPRs.....2
- New Foley Catheter Trays.....2
- Quantiferon-TB Gold Test.....2
- Safe Infusion Practice for Spiking IV Bags.....2
- Stool Specimens.....2
- TJC Survey Prep - Food and Beverages in Patient Care Areas.....2

## Droplet Precautions - Personal Protective Equipment (PPE)


★ Reminder... For patients on Droplet Precautions a new surgical mask should be donned when caring for patients on isolation. A new mask must be donned each time when entering a patient room.

Ear loop and tie masks are appropriate, either may be worn.

There is not a shortage of surgical masks. Please monitor supplies throughout the day and submit requests for re-stocking before supplies are exhausted.



## Ebola



Signage has been revised and will be available within the next few weeks.

## Stool Specimens

New PCR technology is now used for testing stool specimens for bacteria, viruses, and parasites. The testing is not impacted if stool specimens are mixed with urine. Therefore if stool is accidentally mixed with urine, the lab will still accept this specimen. Please do not delay collection of stool specimen.

## Disposable Wipes



\*Don't rush to flush! Disposable wipes used for patient care and environmental cleaning are NOT to be flushed in the toilet. Do not put wipes into toilets, discard in trashcan. Please take every opportunity to educate visitors and family to refrain from disposing these items into the toilet.

## Collection Technique for Quantiferon-TB Gold Test

Please follow instructions when collecting specimens for the QFT, **gently invert tube, do not shake**. Not following the correct collection technique can lead to repeat testing and an unnecessary delay.

## Linen Hampers Required for Contact Isolation

For patients on isolation, please request a yellow disposable hamper to be placed in the patient room for disposition of linen. Do not place linen on floor.

## Safe Infusion Practice for Spiking IV Bags

IV bags should be spiked as close to the time of administration as possible, not to exceed one hour in advance of their use. The only exceptions to this are Code Red Rooms, Emergent Open Heart and ECMO cases. These bags must be dated and timed for 24 hours.

## Fit Test Products

Select fit test products are now available for individual purchase in SDS. These products include the sensitive bitter solution, bitter solution, and hoods.

## The Joint Commission Survey Prep - Food and Beverages in Patient Care Areas

Just a friendly reminder, it is an OSHA violation to have food and beverages in patient care areas.



## New Foley Catheter Trays

To assist in the reduction catheter-associated urinary tract infections, we will be implementing a new foley catheter tray. The new trays are expected to be in circulation by the end of March.

### Infection Control and Prevention

1251 S. Cedar Crest Blvd., Suite 204  
Allentown, PA 18103

2545 Schoenersville Rd., 4th Floor  
Bethlehem, PA 18017

### Brought to you by the Infection Control and Prevention Department

Alida Ammon • Terry Burger • Paula Cook • Carolyn Cramsey • Deborah Fry  
Bernadette Kratzer • Bethann Kretzman • Amber Kucher • Michelle Maron • Sharon Miller  
Sarah Rinker • Robin Tannous • Gary Tallarita • Elaine Walz • Diane Weslosky

A PASSION FOR BETTER MEDICINE

