Lehigh Valley Health Network LVHN Scholarly Works

Department of Pharmacy

Retrospective Study to Identify Predictors of Withdrawal in Pediatric Intensive Care Patients on Long-Term Sedation

Lindsay Fakete PharmD Lehigh Valley Health Network, Lindsay.Fakete@lvhn.org

Jenny Boucher PharmD, JD, BCPS Lehigh Valley Health Network, Jenny.Boucher@lvhn.org

Scott Wheatley MD Lehigh Valley Health Network, Scott_M.Wheatley@lvhn.org

Follow this and additional works at: https://scholarlyworks.lvhn.org/pharmacy

Part of the Pediatrics Commons, and the Pharmacy and Pharmaceutical Sciences Commons Let us know how access to this document benefits you

Published In/Presented At

Fakete, L., Boucher, J. & Wheatley, S. (2014, December, 6). *Retrospective Study to Identify Predictors of Withdrawal in Pediatric Intensive Care Patients on Long-Term Sedation.* Poster session presented at the UHC and ASHP Midyear Conferences, Sacramento, CA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Retrospective Study to Identify Predictors of Withdrawal in Pediatric Intensive Care Patients on Long-Term Sedation Lindsay Fakete, PharmD; Jenny Boucher, PharmD, JD, BCPS; and Scott Wheatley, MD Lehigh Valley Health Network, Allentown, PA

Purpose

The objective of this retrospective study is to examine elements of continuous sedation to determine if they are significant predictors of withdrawal in patients in the pediatric intensive care unit (PICU). This study will also evaluate the documentation of withdrawal symptoms, current weaning approach patterns in the PICU, and compare the approaches with national clinical guidelines for utilization of a sedation wean in this population.

Background

- Critically ill children routinely receive sedative medications in doses and over periods of time that can lead to tolerance and dependence.¹
- Incidence of withdrawal symptoms has been shown to be related to duration of continuous sedation as well as dosage of medications used.^{1,2}
 - Patients who receive 7-14 days of continuous sedation may not require weaning.³
 - Patients who receive more than 14 days will usually need a wean.³
- Withdrawal symptoms can lead to negative consequences such as increased morbidity, higher costs of hospitalization due to longer stays in the intensive care unit, and psychological distress.^{2,4}
 - Typical withdrawal symptoms include agitation, restlessness, irritability, tremors, inconsolable crying, tachypnea, vomiting, diarrhea, and many others.¹⁻⁵

Study Design

IRB approved retrospective medical record review Inclusion Criteria

- June 30, 2014
- without interruption

Exclusion Criteria

- Patients 18 years of age or older
- Patients younger than 2 months of age
- their home medication list
- Patients with a documented diagnosis of diffuse axonal injury
- predictors of withdrawal in patients in the PICU on continuous sedation for at least 7 days.
- The secondary outcomes of the study will be to report:
 - - 7 to 14 days
 - Greater than 14 days
 - - Frequency of dosing
 - Duration of wean
 - Utilization of as needed doses

Disclosure:

Authors of this presentation have the following to disclose concerning possible financial or personal relationships with commercial entities that may have a direct or indirect interest in the subject matter of this presentation.

- Lindsay Fakete nothing to disclose Jenny Boucher – nothing to disclose
- Scott Wheatley nothing to disclose

- Patients between 2 months and 18 years of age admitted to the LVHN PICU between July 1, 2009 and

- Patients must have required continuous sedation of fentanyl and/or midazolam for at least seven days

- Patients who were receiving opioids and/or benzodiazepines prior to admission to the PICU according to

- Patients with a documented diagnosis of autonomic dysfunction syndrome during the admission

• The primary outcome of this study will be to determine if length of continuous sedation, presence of wean from sedation (utilizing methadone and/or lorazepam), and total dose exposed (mg/kg) are significant

- Percent of patients weaned from continuous sedation based on length of sedation split into categories

- To further characterize the approach utilized to wean patients from continuous sedation based on:

Methods

• PICU admission records will be used to generate a list of patients • Variables of interest include:

- Utilization of a wean from sedative medications
- Use of methadone and lorazepam to prevent withdrawal symptoms
- Amount of time patients received continuous sedation
- Total dose of sedation to which the patient was exposed

• Then the presence or absence of withdrawal symptoms will be evaluated in patients:

- If they received a rescue dose of fentanyl, midazolam, morphine, methadone or lorazepam within 72 hours after discontinuation of sedation **and/or**
- If withdrawal symptoms were documented in the medical record by the nurse or physician

Additional data will be collected to describe the study sample and the various weaning techniques utilized

References:

Anand KJS, Willson DF, Berger J, et al. Tolerance and withdrawal from prolonged opioid use in critically ill children. *Pediatrics*. 2010.125: e1208-e1225.

2 Fisher D, Grap MJ, Younger JB, et al. Opioid withdrawal signs and symptoms in children: frequency and determinants. Heart Lung. 2013:407-413.

3 Galinkin J, Koh JL. Recognition and management of iatrogenically induced opioid dependence and withdrawal in children. Pediatrics. 2014; 133: 152-155.

4 Birchley G. Opioid and benzodiazepine withdrawal syndromes in the paediatric intensive care unit: a review of recent literature. Nurs Crit Care. 2009. 14(1): 26-37.

5 Jeffries SA, McGloin R, Pitfield AF, et al. Use of methadone for prevention of opioid withdrawal in critically ill children. Can J Hosp Pharm. 2012; 65(1):12-18.

© 2014 Lehigh Valley Health Network

A PASSION FOR BETTER MEDICINE.



610-402-CARE LVHN.org