

Comparison of Pediatric Outpatient Antibiotic Prescribing Patterns for Specified Indications in Primary Care Practices

Samantha Spishock PharmD

Lehigh Valley Health Network, Samantha.Spishock@lvhn.org

Kristen Held Wheatley PharmD, BCOP

Lehigh Valley Health Network, Kristin_M.Held@lvhn.org

Tibisay Villalobos MD, FAAP

Lehigh Valley Health Network, tibisay.villalobos@lvhn.org

Follow this and additional works at: <https://scholarlyworks.lvhn.org/pharmacy>



Part of the [Medical Education Commons](#), and the [Pharmacy and Pharmaceutical Sciences Commons](#)

Published In/Presented At

Spishock, S., Held Wheatley, K., Villalobos-Fry, T. (2015, December 4). *Comparison of Pediatric Outpatient Antibiotic Prescribing Patterns for Specified Indications in Primary Care Practices*. Poster presented at: Midyear Clinical Meeting; American Society of Health-System Pharmacists, New Orleans, LA.

Spishock, S., Held Wheatley, K., Villalobos-Fry, T. (2016, May 2-4). *Comparison of Pediatric Outpatient Antibiotic Prescribing Patterns for Specified Indications in Primary Care Practices*. Poster Presented at: The Eastern States Conference for Pharmacy Residents and Preceptors, Hershey, PA.

Comparison of Pediatric Outpatient Antibiotic Prescribing Patterns for Specified Indications in Primary Care Practices

Samantha Spishock, PharmD; Kristin Held Wheatley, PharmD, BCOP; Tibusay Villalobos-Fry, MD, FAAP
Lehigh Valley Health Network, Allentown, Pennsylvania

PURPOSE

The objective of this study is to evaluate antibiotic utilization for pediatric patients in the outpatient setting and to describe characteristics related to the patient, prescriber, or specified conditions that may lead to increased antibiotic prescribing.

BACKGROUND

- Antibiotics are the most commonly prescribed medication class in pediatrics, with the use of broad-spectrum antibiotics for pharyngitis on the rise.^{1,2}
- Efforts to decrease antibiotic use for conditions where they are not indicated in the pediatric population and research on this subject have been mainly focused on the inpatient setting.¹
 - Guidelines written by the Infectious Diseases Society of America acknowledge the importance of improving outpatient antibiotic use, but do not specify interventions or recommendations on the implementation of an outpatient antimicrobial stewardship program due to a lack of data in this area.³
- An outpatient antimicrobial stewardship intervention, consisting in part of clinician education, has been shown to significantly reduce off-guideline antibiotic use.⁴
 - Decreasing guideline-discordant broad-spectrum antibiotic use is an area where improvement is necessary, as broad-spectrum third-generation cephalosporins are often utilized for conditions where they are not indicated in the pediatric outpatient setting.⁵
- Provider specialty can also be a factor in increased antibiotic prescribing.
 - Past studies have shown that family medicine providers are more likely to prescribe antibiotics for conditions not requiring antibiotic therapy than general internal medicine providers or pediatric specialists.^{6,7}
- The results of this study may be utilized to implement a clinical education program aimed to improve outpatient antimicrobial utilization in the pediatric population.

STUDY DESIGN

- Retrospective chart review of outpatient encounters
- Inclusion criteria
 - Age 1 month to less than 18 years
 - Outpatient encounter for diagnosis of upper respiratory tract infection (including common cold), bronchitis, bronchiolitis, and/or pharyngitis between March 2, 2015 and September 1, 2015
- Exclusion criteria
 - Neonates <30 days postnatal age
 - Other identified bacterial infection or ongoing bacterial infection coded during same outpatient encounter as above diagnoses
 - Patients with complex chronic conditions⁸
- The primary outcome of this study will be to calculate the percentage of encounters which resulted in an antibiotic prescription out of all encounters for the conditions specified for which an antibiotic is not indicated.

References:

1. Saleh EA, Schroeder DR, Hanson AC, Banerjee R. Guideline-concordant antibiotic prescribing for pediatric outpatients with otitis media, community-acquired pneumonia, and skin and soft tissue infections in a large multispecialty healthcare system. *Clin Res Infect Dis*. 2015;2(1). pii: 1010.
2. Dooling KL, Shapiro DJ, Van Beneden C, Hersh AL, Hicks LA. Overprescribing and inappropriate antibiotic selection for children with pharyngitis in the United States, 1997-2010. *JAMA Pediatr*. 2014;168:1073-4.
3. Dellit TH, Owens RC, McGowan JE Jr, et al; Infectious Diseases Society of America; Society for Healthcare Epidemiology of America. Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America guidelines for developing an institutional program to enhance antimicrobial stewardship. *Clin Infect Dis*. 2007;44:159-77.
4. Gerber JS, Prasad PA, Fiks AG, Localio AR, Grundmeier RW, Bell LM, Wasserman RC, Keren R, Zaoutis TE. Effect of an outpatient antimicrobial stewardship intervention on broad-spectrum antibiotic prescribing by primary care pediatricians: a randomized trial. *JAMA*. 2013;309:2345-52.
5. Vaz LE, Kleinman KP, Raebel MA, Nordin JD, Lakoma MD, Dutta-Linn MM, Finkelstein JA. Recent trends in outpatient antibiotic use in children. *Pediatrics*. 2014;133:375-85.
6. Barlam TF, Morgan JR, Wetzler LM, Christiansen CL, Drainoni ML. Antibiotics for respiratory tract infections: a comparison of prescribing in an outpatient setting. *Infect Control Hosp Epidemiol*. 2015;36:153-9.
7. Hicks LA, Bartoces MG, Roberts RM, Suda KJ, Hunkler RJ, Taylor TH Jr, Schrag SJ. US outpatient antibiotic prescribing variation according to geography, patient population, and provider specialty in 2011. *Clin Infect Dis*. 2015;60:1308-16.
8. Feudtner C, Christakis DA, Connell FA. Pediatric deaths attributable to complex chronic conditions: a population-based study of Washington State, 1980-1997. *Pediatrics*. 2000;106(1 Pt 2):205-9.
9. Hall SA, Kaufman JS, Ricketts TC. Defining urban and rural areas in U.S. epidemiologic studies. *J Urban Health*. 2006;83(2):162-175.

Disclosure:

Authors of this presentation have the following to disclose concerning possible financial or personal relationships with commercial entities that may have a direct or indirect interest in the subject matter of this presentation:

- Samantha Spishock – nothing to disclose
- Kristin Held Wheatley – nothing to disclose
- Tibusay Villalobos-Fry – nothing to disclose

METHODS

- Office encounters for specified conditions between March 2, 2015 and September 1, 2015 will be reviewed for pediatric patients within Lehigh Valley Physician Group (LVPG) practices that utilize the electronic health record (EHR), Epic.
- An antibiotic prescription for any of the specified conditions will be considered non-indicated.
- Our goal will be to include at least 60 different providers, each having a minimum of 5-10 encounters for a specified condition for pediatric patients during the designated time period.
- Data to be collected will include:
 - Patient age, gender, and health insurance coverage
 - Provider specialty, practice site, practice location (urban, suburban, or rural),⁹ and level of provider training
 - Date of encounter, diagnosis assigned, and whether or not an antibiotic was prescribed at the encounter
 - Antibiotic class if an antibiotic was prescribed
- The percentage of overall encounters which resulted in an antibiotic being prescribed will be reported. Descriptive statistics will be used to summarize the characteristics of the encounters as a whole, such as patient age, gender, type of insurance coverage, season of the year in which the encounter occurred, the diagnosis, and the type of antibiotic if one was prescribed.
- The percentage of encounters that were with a pediatric provider and a family medicine provider will be reported.

© 2015 Lehigh Valley Health Network

A PASSION FOR BETTER MEDICINE.™

610-402-CARE LVHN.org