#### Lehigh Valley Health Network

#### **LVHN** Scholarly Works

Department of Medicine

#### Case of an Acute Dual Plaque Rupture

Adrian C. Bell DO *Lehigh Valley Health Network*, Adrian\_C.Bell@lvhn.org

Amit N. Nanavati MD Lehigh Valley Health Network, Amit\_N.Nanavati@lvhn.org

Sarang S. Mangalmurti MD

Lehigh Valley Health Network, Sarang\_S.Mangalmurti@lvhn.org

Follow this and additional works at: https://scholarlyworks.lvhn.org/medicine

Part of the Cardiology Commons, and the Medical Sciences Commons

#### Let us know how access to this document benefits you

#### Published In/Presented At

Bell, A., Nanavati, A., & Mangalmurti, S. (2012). Case of an acute dual plaque rupture. Poster presentation.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

# Case of an Acute Dual Plaque Rupture

Adrian Bell DO, Amit Nanavati MD, Sarang Mangalmurti MD, Division of Cardiovascular Diseases

Lehigh Valley Health Network, Allentown, PA


### Presentation:

- We present a case of a 52 year old gentleman with a history of hypertension who presented to the emergency department with substernal chest pressure and shortness of breath.
- The presenting ECG revealed marked ST-elevation as shown here.

He received the standard aspirin 324 mg, clopidogrel 600 mg,

and a 5,000 unit IV heparin bolus and taken emergently to the


Treatment Course:



### Catheterization:

 Because of the ongoing hemodynamic instability, the left coronary system was engaged prior to the stent being placed which revealed a 100% occlusion of the proximal LAD.

Proximal LAD Occlusion



LAD occlusion in the RAO caudal view

### Proximal LAD Occlusion



Proximal LAD occlusion in the LAO caudal view

### He was taken to the cardiac ICU in critical condition and discharged from the hospital 1 week later with an external defibrillator for severely depressed LV function and frequent ventricular ectopy.



 LAD after placement of stent with restoration of flow




RCA after placement of stent showing restoration of flow

# Catheterization:


catheterization lab.

- A right femoral artery approach was utilized.
- Here, vasopressors were initiated for profound hypotension. The RCA was engaged first based on the presenting ECG and revealed a 100% proximal occlusion.
- During the intervention, multiple episodes of ventricular tachycardia/fibrillation occurred requiring ACLS and intubation.





LV Gram



Left ventriculogram revealing severe anterior and inferior wall akinesis

## Treatment:

- Both the proximal RCA and proximal LAD were successfully reperfused and a bare-metal stent was placed in the RCA and a drug-eluting stent placed to the proximal LAD.
- Both interventions led to a 0% residual stenosis with resultant TIMI 3 flow.

### Discussion:

- A simultaneous acute dual-plaque rupture resulting in two coronary arteries being occluded is a rare occurrence. This case demonstrates that proper and timely visualization of all the coronary arteries during an acute myocardial infarction is critical. During an acute myocardial infarction, time to vessel reperfusion is paramount and a delay to visualize other vessels can cause a delay to reperfusion.
- One option is to visualize the presumed non-infarct artery while a balloon and stent are being set-up. This can be done quickly and accurately and would allow for proper decision making. This case demonstrates the importance of the visualization of all the coronary arteries during an acute myocardial infarction.

A PASSION FOR BETTER MEDICINE."

