

Interrupted Aortic Arch in the Adult

Amit Nanavati MD
Lehigh Valley Health Network

Matthew W. Martinez
Lehigh Valley Health Network, matthew_w.martinez@lvhn.org

Nainesh Patel MD
Lehigh Valley Health Network, nainesh_c.patel@lvhn.org

Follow this and additional works at: <https://scholarlyworks.lvhn.org/medicine>



Part of the [Cardiology Commons](#), and the [Medical Sciences Commons](#)

Let us know how access to this document benefits you

Published In/Presented At

Nanavati, A., Martinez, M., & Patel, N. (2013). *Interrupted aortic arch in the adult*. Poster presentation.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Interrupted Aortic Arch in the Adult

Amit Nanavati, MD; Matthew Martinez, MD; Nainesh Patel, MD; Division of Cardiology

Lehigh Valley Health Network, Allentown, Pennsylvania

Introduction:

- Interrupted aortic arch is a rare phenomena, occurring in approximately 3 of 1,000,000 cases. It was first recognized in 1778.¹
- Survival to adulthood is an even more rare occurrence with 30 cases reported.³
- Three types have been described which describe the relation of the interruption to the great vessels.^{1,3}
 - Type A – distal to left subclavian artery (43%)
 - Type B – between left carotid and subclavian arteries (53%)
 - Type C – between the brachiocephalic and left carotid arteries (4%)
 - Of note, other studies³ have reported Type A to be the most common

Case Presentation:

We describe a 53 year old male who presented for cardiac catheterization for pre-operative planning for severe aortic insufficiency (presumably from a history of rheumatic disease). He originally presented for several months of worsening dyspnea with exertion.

Physical Examination: Physical exam revealed >30 mmHg difference in systolic pressure between the upper extremities and lower extremities bilaterally without significantly different pressures in the left and right arms. (Figure 1)

Chest X-Ray: Demonstrated cardiomegaly. (Figure 2).

Echocardiography revealed a tricuspid aortic valve with moderate to severe regurgitation and severe dilation of the ascending aorta.

Aortography: Aortography via a femoral approach revealed an interrupted arch distal to the left subclavian artery with large subcostal arteries serving as “bridging” collateral vessels. (Figure 3)

CT Angiography: CT Angiography demonstrated significant enlargement of the intercostal arteries as well as collateralization from the internal mammary arteries to the inferior epigastric arteries. A fibrous strand connected the ascending and descending aortas distal to the left subclavian artery. (Figures 4 & 5)

References:

- 1 Messner et. al. Interrupted Aortic Arch in the Adult. Texas Heart Institute J. 2002; 29(2): 118-121.
- 2 Ogino et. al. Two stage repair for aortic regurgitation with interrupted aortic arch. Ann Thor Surg 1998 Apr; 65(4): 1151-3.
- 3 Borgohain et. al. Isolated Interrupted Aortic Arch in an 18-Year Old Man. Texas Heart Institute J. 2013; 40(1): 79-81.
- 4 Sai Krishna et. al. Interruption of Aortic Arch in Adults. Texas Heart Institute J. 2005; 32(2): 147-150.




Figure 1: Schematic of Blood Pressure and Pulse Recordings.

Figure 2: Chest x-ray demonstrating cardiomegaly.

Figure 3: Aortic angiogram demonstrating interruption and collateral vessels (arrows).

Figure 4: Interruption in the aortic arch (arrow) is noted distal to the left subclavian artery.

Figure 5: Significant collateralization is noted from the internal mammary to the inferior epigastric arteries (arrow). Enlarged intercostal arteries were also noted.



Discussion:

- Surgical intervention is high-risk in this patient due to the difficulty in achieving hemostasis due to massively enlarged intercostal arteries.
- However, he subsequently underwent unremarkable aortic valve replacement with plans for future aortic arch replacement.
- Previous surgical approaches have included 1 or 2 stage repairs¹, though two stage repair has been advocated in one study with repair of the aortic valve.²
- Consideration should also be given to extra-anatomic bypass³ of which ventral aortic repair has been shown to have good outcomes.⁴
- Interrupted aortic arch is a rare condition, however association with aortic arch dilation and aortic insufficiency seems to be even less frequent.

© 2013 Lehigh Valley Health Network

A PASSION FOR BETTER MEDICINE.™

Lehigh Valley Health Network

610-402-CARE LVHN.org