

Visualization and Management of a Distally Embolized Stent After ST-Elevation Myocardial Infarction

Amit Nanavati MD
Lehigh Valley Health Network

Nainesh C. Patel MD
Lehigh Valley Health Network, nainesh_c.patel@lvhn.org

Follow this and additional works at: <https://scholarlyworks.lvhn.org/medicine>



Part of the [Cardiology Commons](#), and the [Medical Sciences Commons](#)

Published In/Presented At

Nanavati, A., & Patel, N. (2013). *Visualization and management of a distally embolized stent after ST-Elevation myocardial infarction*. Poster presentation.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Visualization and Management of a Distally Embolized Stent After ST-Elevation Myocardial Infarction

Amit Nanavati, MD; Nainesh Patel, MD
Lehigh Valley Health Network, Allentown, Pennsylvania

Background:

- Stent embolization is a rare complication in the current era of percutaneous intervention with an incidence as low as 0.3%.¹
- Common causes of this phenomena include dislodgement from the balloon, direct stenting as well as attempting to cross a calcified lesion.²
- Previous methods for removing the embolized stent have included snares, balloons and distal protection devices.³
- “Crush” stenting as well as conservative management have also been advocated.
- Visualization can also be a challenge as angiography and intravascular ultrasound have inherent limitations.

Case Presentation:

- 63 year-old male with a history of tobacco abuse, morbid obesity and hypertension who presented with 4 hours of substernal chest pressure.
- Initial electrocardiogram demonstrated ST-elevation in the inferior leads and he was taken for emergent coronary angiography.
- A severe lesion was seen in the proximal RCA (**Figure 1**) and a Veriflex 5.0 x 16 mm bare-metal stent was placed in the proximal RCA with no angiographically apparent residual stenosis. (**Figure 2**).
- A second stent placement was also attempted in the proximal RCA, however, visualization was noted to be suboptimal due to body habitus.
- Later that day, the patient had recurrent symptoms of chest pain and inferior lead ST-elevations.
- Repeat coronary angiography (**Figure 3**) with the assistance of optical coherence tomography noted that the second RCA stent was partially deployed and had embolized into the PDA. (**Figures 4-5**).

Results:

- The stent was then crushed against the vessel wall with a 2.75 x 23 mm Xience Everolimus eluting stent.
- The embolized stent could not be fully covered with the new stent.
- Optical coherence tomography also visualized a 60% lesion in the deployed RCA stent, which was restented.

Conclusion:

- The patient was discharged two days later in stable condition.
- This case illustrates that optical coherence tomography is a useful tool in visualizing distally embolized stents when limitations make other methods impractical.




Figure 1 — Initial severe stenosis in the proximal RCA.




Figure 2 — No apparent residual stenosis in the RCA.




Figure 3 — Angiography demonstrating likely location of stent embolization in the PDA (arrow).




Figure 4 — Optical coherence tomography depicting embolized stent.




Figure 5 — The full extent of the embolized stent could not be covered.

© 2013 Lehigh Valley Health Network

References:

- 1 Saleh L. et al. Exp Clin Cardiol 2010; 15(3): e70-2.
- 2 Farman M. et al. J Pak Med Assoc 2010; 60(2): 140-2.
- 3 Guiguari P et. al. J Invasive Card 2005; 17(3).

A PASSION FOR BETTER MEDICINE.™

Lehigh Valley
Health Network

610-402-CARE LVHN.org