Lehigh Valley Health Network

LVHN Scholarly Works

Research Scholars Poster Presentation

Potential Extracorporeal Membrane Oxygenation Use for Increased Survival of In-Hospital Cardiac Arrests

Amanda Broderick

James K. Wu MD

Lehigh Valley Health Network, james.wu@lvhn.org

Heather Geist heather.geist@lvhn.org

Follow this and additional works at: https://scholarlyworks.lvhn.org/research-scholars-posters
Let us know how access to this document benefits you

Published In/Presented At

Broderick, A., Geist, H., Wu, J., (2014, July, 25) *Two- Year Retrospective Study of In-Hospital Cardiac Arrests.* Poster presented at LVHN Research Scholar Program Poster Session, Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Potential Extracorporeal Membrane Oxygenation Use for Increased Survival of In-Hospital Cardiac Arrests

Amanda Broderick, James Wu, M.D., Heather Geist Department of Surgery, Division of Cardiothoracic Surgery Lehigh Valley Health Network, Allentown, Pennsylvania

Background

In-hospital cardiac arrest (IHCA) is a relatively uncommon event but is associated with a low rate of survival.

Extracorporeal membrane oxygenation (ECMO) is a highly aggressive and advanced cardiopulmonary support system that can benefit select patients who cannot be successfully resuscitated through traditional cardiopulmonary resuscitation (CPR). It temporarily allows for adequate bodily perfusion in patients in refractory cardiac arrest and allows for valuable time so that the pathologies behind the cardiac arrests can be evaluated and treated.¹

Methods

A retrospective study was done involving 182 patients in 2011 and 2012 within the Lehigh Valley Health Network for which a "code blue" was called for an in-hospital cardiac arrest.


Data was examined for each patient from electronic medical and resuscitation records. The overall exclusion criteria was an age <18 or >70 years or an existing Do Not Resuscitate (DNR) before coding.


REFERENCES

1 Lazzeri C., Bernardo P., Sori A., Innocenti L., Stefano P., Peris A., gensini G.F., Valente S. Venous-arterial extracorporeal membrane oxygenation for refractory cardiac arrest: a clinical challenge. *Eur Heart J Acute Cardiovasc Care* 2013; 2: 118-26.


2 Haneya A., Philipp A., Diez C., Schopka S., Bein T., Zimmermann M., Lubnow M., Luchner A., Agha A., Hilker M., Hirt S., Schmid C., Muller T. A 5-year experience with cardiopulmonary resuscitation using extracorporeal life support in non-postcardiotomy patients with cardiac arrest. *Resuscitation* 2012; 83: 1331-37.

Results & Figures










Results



Of the 131 patients that could not be permanently resuscitated, 44 would have been good candidates for ECMO-CPR. Previous studies have found survival rates for patients undergoing ECMO-CPR due to IHCA to be 30-42%. This indicates that approximately 13-18 additional patients could have survived at LVHN in 2011-2012, increasing the overall IHCA survival rate from 28% to 37%.

Conclusions

Introducing ECMO-CPR for select IHCA patients in the Lehigh Valley Health Network is a realistic way to produce positive outcomes for patients who would otherwise face grave prognoses and better the overall survival statistics for IHCA.

© 2014 Lehigh Valley Health Network



Special thanks to Brandon Follick for his help!