Lehigh Valley Health Network

LVHN Scholarly Works

Research Scholars Poster Presentation

Surgical Treatment of Multiple Vein Graft Aneurysms in 72 Year Old Man

Heather Geist heather.geist@lvhn.org

James K. Wu MD

Lehigh Valley Health Network, james.wu@lvhn.org

Raymond Singer MD Lehigh Valley Health Network, raymond.singer@lvhn.org

Follow this and additional works at: https://scholarlyworks.lvhn.org/research-scholars-posters

Let us know how access to this document benefits you

Published In/Presented At

Geist, H., Wu, J., Singer, R., (2014, July, 25) *Surgical Treatment of Multiple Vein Graft Aneurysms in a 72 Year Old Man.* Poster presented at LVHN Research Scholar Program Poster Session, Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Surgical Treatment of Multiple Vein Graft Aneurysms in 72 Year Old Man

Heather Geist, Raymond Singer MD, James Wu MD

Department of Surgery, Division of Cardiothoracic Surgery, Lehigh Valley Health Network, Allentown, Pennsylvania

Introduction

Saphenous vein graft (SVG) aneurysms following coronary artery bypass graft (CABG) surgery are rare but fatal complications that frequently require surgical repair. This study addresses a rare surgical case involving two SVG aneurysms with thrombus in an 72 year old man, requiring a third sternotomy 22 years following CABG.




Figure 1. Aneurysmal right coronary artery bypass graft prior to resection

Methods

All information was gathered retrospectively using the LVHN medical records and database. Patient data was analyzed primarily from 2007 to the present. Surgical dictations, CT scans, and chest x-rays were all reviewed.



Results




Figure 2. Patient CT scan prior to surgery showing the 8 cm RCA graft and 5 cm circumflex graft aneurysms




Figure 3. Patient CT scan following surgery showing the removal of the 8 cm RCA graft aneurysm

Year Preliminary Cause for Surgery Surgical Procedures CABG x3, SVGs to LAD, second Emergent, unstable angina marginal and RCA Significant dyspnea, substernal chest pressure, catheterization showed LAD with Redo sternotomy, CABG x2, LIMA to stenosis, RCA graft occluded and aneurysmal, LAD and a SVG to second marginal second marginal occluded Redo sternotomy, resection of giant Following syncope episode, growth of RCA RCA vein graft aneurysm, CABG x1 with CryoVein to distal posterior graft aneurysm from 4 cm to 8 cm descending artery

Table 1. Significant cardiothoracic surgical procedures and timeline of occurrence with description




Figure 4. Chest x-ray of patient with evident aneurysmal bulge of RCA




Figure 5. Chest x-ray of patient following surgery with removal of RCA aneurysm

Conclusion

True or pseudoaneurysms can arise following CABG surgery. It was determined that this patient presented with two true aneurysms, which are frequently late complications after surgery.² Our patient presented with common symptoms accompanying aneurysms, including a syncope episode,³ history of hyperlipidemia, and a history of myocardial infarction.⁴

A previous study using a small cohort had found that SVG aneurysms to the RCA and circumflex were two of the three most frequent aneurysmal grafts.⁴ Complications experienced by this patient as a result of the aneurysm lead to necessary surgical intervention.

References

¹Drummer E, Furey K, Hollmann J. Rupture of a saphenous vein bypass graft during coronary angioplasty. *Br Heart J.* 1987;**58**:78-81.

²Liang BT, Antman EM, Taus R, *et al.* Atherosclerotic aneurysms of aortocoronary vein grafts. *Am J Cardio.* 1988;**61**:185-188.

³Mayglothling J, Thomas MP, Nyzio JB, *et al.*Aneurysm of aortocoronary saphenous vein graft: case report and literature review. *Heart Surg Forum*. 2004;**55**:587-588.

⁴Sareyyupoglu, B, Schaff HV, Ucar I, *et al.* Surgical treatment of saphenous vein graft aneurysms after coronary artery revascularization. *Ann Thorac Surg.* 2009;**88**:1801-1805.