Lehigh Valley Health Network LVHN Scholarly Works

Research Scholars Poster Presentation

Use of Thrombolytic Therapy for Complicated Pleural Effusions

Lauren E. Sterner Lauren.Sterner@lvhn.org

Kelly M. Wanamaker MD Lehigh Valley Health Network, Kelly.Wanamaker@lvhn.org

Katherine M. Penetar DNP, CRNP Lehigh Valley Health Network, Katherine_M.Penetar@lvhn.org

Sanjay M. Mehta MD Lehigh Valley Health Network, Sanjay_M.Mehta@lvhn.org

Follow this and additional works at: https://scholarlyworks.lvhn.org/research-scholars-posters Let us know how access to this document benefits you

Published In/Presented At

Sterner, L. Wanamaker, K. Penetar, K. Mehta, S. (2019, August). *Use of Thrombolytic Therapy for Complicated Pleural Effusions.* Poster Presented at: LVHN Research Scholar Program Poster Session, Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Use of Thrombolytic Therapy for Complicated Pleural Effusions

BACKGROUND

- A complicated pleural effusion is defined as an accumulation of pleural fluid with fibrin deposits and a non-elastic pleural membrane peel.¹
- Treatment remains controversial; traditionally, surgical decortication has been the mainstay, however there are contraindications and an associated high morbidity.²
- Intrapleural thrombolytic therapy offers an alternative to surgical intervention and alteplase has been shown to be an effective treatment.³

Primary Objective: To determine the efficacy of intrapleural alteplase therapy in treatment of complex pleural effusion. Successful treatment is defined as resolutions or near resolutions in chest radiographs. **Secondary Objective**: To compare the length of hospital stay between traditional surgical treatment and that of intrapleural thrombolytic therapy.

METHODS

Retrospective chart review on adult admissions with a complicated pleural effusion treated with either thrombolytic therapy, pigtail tube, or surgical decortication from January 2016 to May 2019.

The following information was collected: age, sex, etiology of pleural effusion, treatment, dosages of alteplase, success of treatments, complications and length of stay.

Descriptive statistics were used to evaluate the primary and secondary objectives and to summarize the details of each encounter.

1. Walker, C., Shirk, M., Tschampel, M., & Visconti, J. (2003). Intrapleural Altepase in a Patient with Complicated Pleural Effusion. The Annals of Pharamcology, 37, 376-379. doi:10.1345/aph.1C248 2. Piccolo, F., Popwicz, N., Wong, D., & Lee, Y. (2015). Intrapleural tissue plasminogen activator and deoxyribonuclease therapy for pleural infection. Journal of Thoracic Disease, 7(6), 999-1008.

doi:10.3978/j.issn.2072-1439.2015.01.30 3. Ben-Or, S., Feins, R., Veeramachaneni, N., & Haithcock, B. (2011). Effectiveness and Risks Associated With Intrapleural Alteplase by Means of Tube Thoracostomy. The Society of Thoracic Surgeons, 4th ser., 860-864. doi:10.1016/j.athoracsur.2010.10.082

© 2019 Lehigh Valley Health Network

Lauren Sterner, Kelly M. Wanamaker, MD, Katherine M. Penetar DNP, CRNP, Sanjay M. Mehta, MD Department of Cardiothoracic Surgery Lehigh Valley Health Network, Allentown, Pennsylvania

OUTCOMES

- 98 patients total; 70 male, 28 female.
- and resulted in full resolution in 84.4% of cases (27 out of 32 patients).
- Five patients had an incomplete response to alteplase treatment, four of which required videoassisted thoracic surgery (VATS). The one patient was not a surgical candidate and received antibiotic therapy.

Figure 1. Pleural effusion etiology for those with full resolution after thrombolytic therapy treatment.

Parapneumonic

Empyema

44%

15%

Figure 2. Average doses of alteplase based off the outcome of the thrombolytic therapy.

Surgical Intervention

Incomplete Resolution

Complete Resolution 2.0 ± 0.817 2.2 ± 0.837

• 32 (32.7%) were treated with thrombolytic therapy

Loculated

41%

Table 1. Compariso Therapy to Surgio

Incidence of Full Rese Alteplase therapy Surgical interven Complications Alteplase therapy Bleeding Sepsis Respirato Other Surgical intervent Bleeding Sepsis Respirato Other

Length of stay Alteplase therap Surgical interver

- avoid surgery.
- on imaging.

LVHN.org

RESULTS

n of Thrombolytic cal Intervention.	N (%) Md (IQR)
olution	
/	27 (84.4%)
tion	32 (96.7%)
/	
	0 (0%)
	2 (6.25%)
ory failure	2 (12.5%)
	4 (6.25)
tion	
	1 (3.03%)
	5 (15.2%)
ory failure	2 (3.03%)
	1 (3.03%)
У	6.5 (9.5)
ntion	8 (9)

CONCLUSIONS

Overall, surgical intervention resulted in more cases of complete resolution, but had a longer length of stay. Those who had incomplete resolution of the effusion after thrombolytic therapy received less dosages of alteplase and dornase compared to those that did.

• Alteplase therapy proves a good treatment option for most cases of complex fluid collection and in those in which surgery is contraindicated or those looking to

3.67 ± 1.85 • Implementing a standard protocol of alteplase and dornase dosing combined with earlier intervention would optimize cases of complete resolution observed

