Lehigh Valley Health Network

LVHN Scholarly Works

Department of Emergency Medicine

Physician Directed Smoking Cessation in the ED: Do Patients Opt-Out?

Ali S. Aziz BS Lehigh Valley Health Network, Ali.Aziz@lvhn.org

Natalie M. Greco MD Lehigh Valley Health Network, natalie.greco@lvhn.org

Timothy J. Batchelor MD Lehigh Valley Health Network, timothy.batchelor@lvhn.org

Andrew H F Miller DO Lehigh Valley Health Network, Andrew.Miller3@lvhn.org

Theodore Doherty DO Lehigh Valley Health Network, Theodore.Doherty@lvhn.org

See next page for additional authors

Follow this and additional works at: https://scholarlyworks.lvhn.org/emergency-medicine



Part of the Emergency Medicine Commons

Published In/Presented At

Aziz, A., Greco, N., Batchelor, T., Miller, A., Doherty, T., Kruklitis, R., Crowley, L., Casey, E., MacLean, E., Trainer, J., & Greenberg, M. (2020, April). Physician Directed Smoking Cessation in the ED: Do Patients Opt-Out? Poster presented at: PACEP Scientific Assembly, Pittsburgh, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Authors Ali S. Aziz BS; Natalie M. Greco MD; Timothy J. Batchelor MD; Andrew H F Miller DO; Theodore Doherty DO; Robert Kruklitis MD, PhD; Lauren Crowley BA; Edward W. Casey RPh, MBA; Elizabeth A. MacLean PharmD, PhD; JoAnn B. Trainer PharmD; and Marna R. Greenberg DO, MPH, FACEP

Physician Directed Smoking Cessation in the ED: Do Patients Opt-Out?

Natalie M. Greco, MD, Ali S. Aziz, BS, Timothy J. Batchelor, MD, Andrew H. F. Miller, DO, Theodore Doherty, DO, Robert J. Kruklitis, MD, PhD, Lauren M. Crowley, BA, Edward W. Casey, RPh, MBA, Elizabeth A. MacLean, PharmD, PhD, JoAnn B. Trainer, PharmD, Marna Rayl Greenberg, DO, MPH

¹Lehigh Valley Hospital and Health Network/University of South Florida Morsani College of Medicine, Lehigh Valley Campus, Allentown, PA, ²Pfizer Pharmaceutical Company, New York, NY

Background/Objective

SBIRT (Screen, Brief Intervention, and Referral for Treatment) interventions for smoking cessation in the ED have been shown to be effective. The goal of this study was to investigate patients' willingness to participate in a smoking cessation intervention known as STIR (Screening, Treatment Initiation, and Referral).

Methods

In this prospective pilot quality improvement program, a convenience sample of smokers at two NE PA hospitals who met inclusion criteria were approached. Criteria included being: English speaking, ≥18yo, discharged, and not critically ill, incapacitated, incarcerated, or known to be currently pregnant. A standardized intervention was provided by residents/attendings to the patient in which the patient could opt-out at any level (to have the intervention, to receive prescriptive advice, or to follow-up with their primary care provider/tobacco treatment program). At the end of the encounter, patients were asked (5-point Likert scale) how important it was to have this conversation with a physician about their smoking behavior.

Results

A total of 106 patients who met inclusion criteria were approached and 99 (93%) agreed to the intervention. Of those that agreed, 50 (50.5%) were male and 49 (49.5%) were female. Their mean age was 42.4 (range 19–85 years old) and 80% reported having a primary care physician. Patients (as indicated) were offered prescriptive advice (nicotine replacement and/or oral medication) and 58

(58.6%) accepted one or both. An indication for prescriptive advice was not present in 6 (6.1%). Seventy-seven (77.8%) patients opted for at least one ambulatory referral for follow-up with primary care or tobacco treatment program. Seventy-nine (79.8%) felt it was important (Likert scale 4 or 5) for the physician to be the team member to have this conversation with them.

Conclusion

In this small ED pilot using the STIR concepts in a patient "opt-out" setting, nearly all smokers chose to participate in the smoking cessation intervention and the vast majority felt it was important for the physician member of the health care team to lead the discussion. Over half of the patients accepted prescriptive advice and more than ¾ agreed to ambulatory referral for follow-up. These findings support a willingness of patients to participate in STIR. Future study regarding their change in smoking behaviors compared to those that receive SBIRT is indicated.

Acknowledgments

This project was funded, in part, by a quality improvement program grant from Pfizer Inc.









