

## A Review of Interstitial Pneumonitis Caused by Elotuzumab Through Case Discussions and Academic Writings.

Ghulam Ghous  
*University of Missouri*

Hafiz Muhammad Hassan Shoukat  
*Premier Health/Wright State University*

Zahid Ijaz Tarar  
*University of Missouri*

Muhammad Usman Zafar  
*Lehigh Valley Health Network, muhammad.zafar@lvhn.org*

Follow this and additional works at: <https://scholarlyworks.lvhn.org/emergency-medicine>



Part of the [Medicine and Health Sciences Commons](#)

Let us know how access to this document benefits you

---

### Published In/Presented At

Ghous, G., Shoukat, H., Tarar, Z. I., & Zafar, M. U. (2020). A Review of Interstitial Pneumonitis Caused by Elotuzumab Through Case Discussions and Academic Writings. *Cureus*, 12(10), e11162. <https://doi.org/10.7759/cureus.11162>

This Article is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact [LibraryServices@lvhn.org](mailto:LibraryServices@lvhn.org).

# A Review of Interstitial Pneumonitis Caused by Elotuzumab Through Case Discussions and Academic Writings

Ghulam Ghous, Hafiz Muhammad Hassan Shoukat, [...], and Muhammad Usman Zafar

## Abstract

Patients with relapsed or refractory multiple myeloma have undergone two or three previous therapies are now being treated with a humanized IgG1 monoclonal antibody elotuzumab (HuLuc63) that targets F7 signalling lymphocytic activation molecule F7 (SLAMF7)- a signalling lymphocytic activation molecule. It is combined with dexamethasone and lenalidomide/pomalidomide for therapy. Adverse effects associated with elotuzumab consists of peripheral neuropathy, fever, constitutional symptoms (fatigue, headache, decreased appetite), and infections. A rare side effect of interstitial lung disease has only been observed in a single case. There are two case studies presented below of hypoxic respiratory failure upon this monoclonal antibody treatment both were successfully treated with steroid therapy. This article brings forth the hypothesis that elotuzumab can cause pneumonitis, and discontinuation of elotuzumab along with high-dose corticosteroids helps reverse the pneumonitis.

**Keywords:** interstitial pneumonitis, elotuzumab, multiple myeloma

## Introduction

Humanized IgG1 monoclonal antibody elotuzumab (HuLuc63) that targets signalling lymphocytic activation molecule F7 (SLAMF7) [1-3]. Multiple myeloma cell death is caused by it through direct activation of natural killer cells targeted at neoplastic plasma cells through Fc receptors (CD16), SLAMF7 pathway and by tagging them for recognition of ADCC (antibody-dependent cellular toxicity). With little to no expression of SLAMF7 protein in normal tissues, elotuzumab selectively kills multiple myeloma cells with minimal off-target effects [4]. In November 2015, food and drug administration (FDA) approved the combination therapy of elotuzumab with lenalidomide and dexamethasone (Elo-Ld) based on ELOQUENT-2 trial on adult patients with one prior therapy [5]. Elotuzumab combined with pomalidomide and dexamethasone (Elo-Pd) has been approved for use in multiple myeloma patients having received two previous therapies, including lenalidomide and PI (protease inhibitor) in 2018 after ELOQUENT-3 trial [6]. There are numerous side effects associated with elotuzumab combination therapy, including lymphopenia, fatigue, diarrhoea, pyrexia, constipation, cough, peripheral neuropathy and nasopharyngitis [5]. Pulmonary toxicity associated with elotuzumab (combined with lenalidomide/pomalidomide and dexamethasone) was barely described in Phase 1-3 clinical trials [5-7]. Only one case report of elotuzumab induced interstitial lung disease exists [8], with a few cases reports on lenalidomide and pomalidomide induced interstitial lung disease [9-14]. This article covers two cases of acute hypoxic respiratory failure upon administration of elotuzumab, and both treated successfully with corticosteroids.

## Case presentation

### Case 1

A 50-year-old male having a history of multiple myeloma (IgG kappa), on weekly elotuzumab infusion for last five weeks (last infusion four days before admission) along with lenalidomide and dexamethasone presented to emergency with four-day symptoms of shortness of breath, fatigue and malaise. He also reported dry cough. He denied having fever, chills, night sweats, chest pain, abdominal pain, nausea, vomiting, diarrhoea, urinary changes and lower extremity swelling. The patient was diagnosed with pulmonary embolism a month back. Compliance of the patient with therapeutic dose enoxaparin was documented. On initial evaluation, the patient was hypoxic requiring 2-3-litres oxygen, later increased to 15 litres high flow nasal cannula, and eventually stabilized on bilevel positive airway pressure (BiPAP). On physical examination, he had a temperature of 101.3 F, and faint crackles (heard bilaterally at bases) on chest examination. Cardiovascular, gastrointestinal, and neurological examinations were normal.

Laboratory tests revealed thrombocytopenia (platelets (PLT)  $16 \times 10^9/L$ ) and anaemia (Hb 7.8 g/dl) consistent with initial baseline values, acute kidney injury (creatinine 1.28 mg/dl), mild pro-BNP elevation (394 pg/ml) with insignificant electrocardiogram abnormality. Viral respiratory pathogens polymerase chain reaction (PCR), COVID-19 PCR, legionella, and streptococcal urine antigens were negative. CT Chest with pulmonary embolism protocol demonstrated the decreased size of previously seen filling defect within right lower lobe segmental artery but new bilateral diffuse ground-glass opacities (Figure 1). Elotuzumab induced pneumonitis, sepsis, pulmonary edema, and multifocal pneumonia were taken as a differential diagnosis. The patient was admitted, blood and urine cultures were sent and started on vancomycin along with meropenem. Normal systolic function (ejection fraction 50%), normal wall motion with no right to left shunt were observed on echocardiogram with bubble study. Continued administration of antibiotics with an observation period of one to two days was suggested by the haematologist along with administration of corticosteroids for possible elotuzumab induced pneumonitis if the patient did not improve under the antibiotics.




Figure 1

Axial contrast-enhanced CT chest (lung window) showing bilateral diffuse ground-glass opacities.

The patient continued to have shortness of breath requiring 10-15-litres oxygen on high flow nasal cannula and intermittent BIPAP over the next 24 hours. He was started on methylprednisolone (125 mg) every six hours after consultation with the haematologist. This brought down the oxygen requirement to 6 litres/min the next day, but the malaise persisted. Antibiotics were stopped since the urine and blood cultures were negative. Significant improvement was observed on the fourth day of intravenous methylprednisolone as the patient ambulated comfortably. He was not requiring any oxygen, was ambulating without difficulty. The patient was discharged with a prescription of prednisone (40 mg) daily with prolonged taper: 40 mg daily for the first two weeks, 10 mg weekly for next three weeks. Marked improvement of the bilateral opacities was observed after three weeks on chest x-ray (Figure 2). After an additional three weeks, CT chest was ordered, but patient decided to discontinue chemotherapy due to adverse effects, low functional status and opted for hospice care.




Figure 2

(A) Chest x-ray shows diffuse bilateral interstitial and peripheral opacities. (B) Improved bilateral patchy and hazy airspace opacities.

### Case 2

A 73-year-old female was admitted with the symptoms of malaise, cough, and shortness of breath on exertion. Past medical history revealed multiple myeloma (stage III light lambda light chain) on weekly elotuzumab for the last nine weeks (last infusion five days ago). Patient endorsed diffuse weakness, chills and sweats for the past couple of days and a 1-day history of cough with progressively worsening shortness of breath. She was uncertain about fever: denied headache, chest pain, abdominal pain, diarrhoea, dysuria. On presentation to the emergency department, the patient was tachycardic and febrile, temperature 101.4 F. Patient was hypoxic and tachypneic while in the emergency department and required 5-liters oxygen by nasal cannula. On examination, she was arousable though lethargic with left-sided chest crackles.

Comprehensive metabolic panel and blood tests revealed mild hypokalemia, chronic anaemia, mild acute kidney injury, and lactic acid (0.8). Unremarkable electrocardiogram and negative troponins were documented (pro b-type natriuretic peptide (pro-BNP) 108 pg/ml). Mild interstitial pulmonary edema was observed on chest x-ray. She tested negative for COVID-19. Venous blood gases displayed a pH of 7.3, PCO<sub>2</sub> of 41, and PO<sub>2</sub> of 39. Blood cultures were sent to the laboratory for evaluation. The patient was admitted to the hospital with the administration of 1.5 L saline bolus and started on vancomycin and piperacillin/tazobactam. Elotuzumab induced pneumonitis, sepsis, pulmonary edema, and pneumonia were considered under differential diagnosis. Corticosteroid administration was recommended by the haematologist if the patient did not improve under the antibiotics during the observation period of one to two days. CT pulmonary angiogram was negative for pulmonary embolism but showed patchy multifocal, right greater than left ground-glass and reticular airspace opacities (Figure 3). Normal systolic function (ejection fraction 65%) and unremarkable regional wall motion were observed on echocardiogram. Respiratory viral pathogen panel by PCR, urine legionella and streptococcus pneumonia antigen were negative.




Figure 3

(A) Contrast-enhanced CT chest axial section (lung window) demonstrating patchy bilateral, right greater than left reticular and ground-glass airspace opacities, and (B) similar bilateral ground-glass opacities (coronal view)

Patient continued to be tachycardic and tachypneic, requiring oxygen for the next 72 hours while on antibiotics. Blood cultures, sputum cultures and urine cultures were negative. The patient was started on Prednisone 1mg/kg daily on the fourth day of admission, and the antibiotics were stopped. She was down to 2 litres/min oxygen the next day, and the dyspnea improved. On the third day, marked improvement in the cough was observed as ambulation came close to normal. For the next five days, the patient was continued on prednisone (70 mg) and then discharged with prednisone (60 mg) to wean by 10 mg every five days. CT chest was ordered after completion of steroids, but patient decided to discontinue chemotherapy due to adverse effects and opted for hospice care. Her breathing status remained normal for the next four weeks during follow-ups.

## Discussion

Elotuzumab is the first and only immune-stimulatory monoclonal antibody indicated for the combination treatment of patients with multiple myeloma. Mechanism of actions involves targeting SLAMF7. SLAMF7 expression is highest on plasma cells (malignant and normal), natural killer cells, and a



histological and radiological patterns vary, including organizing/hypersensitivity/(non-specific) interstitial pneumonitis, acute respiratory distress syndrome, and diffuse alveolar haemorrhage. Four prior acute lung toxicity reports exist related to pomalidomide [12-14]. It usually takes 8 - 120 days from initiation of treatment to onset of symptoms. Symptoms of immunomodulatory drugs lung toxicity include cough, dyspnea, fever, and hypoxia; ground-glass opacities being the most common radiological findings [12]. Significant take-away from the first case is that despite his long history of treatment with lenalidomide, interstitial pneumonitis developed acutely after elotuzumab was added to the combined therapy. In the second case presented the patient was receiving weekly elotuzumab with pomalidomide 4 mg daily (day 0-21) but stopped taking pomalidomide after 14 days due to fatigue, weakness, loss of appetite and failure to thrive. There were no respiratory symptoms during pomalidomide treatment and immediately after stopping it. It was after nine weeks of weekly administration of elotuzumab that the patient developed acute hypoxic respiratory failure, and we believe her respiratory failure was secondary to elotuzumab and not pomalidomide.

Tanaka et al. suggest that the mechanism by which elotuzumab induces lung injury (mostly unknown) might be linked to TNF- $\alpha$  secretion, as other studies have revealed monoclonal antibodies (e.g., Rituximab) associated ILD is related to TNF- $\alpha$  secretion [8, 16]. Elotuzumab has been demonstrated to induce TNF- $\alpha$  secretion in vitro [17]. It has also shown a synergistic effect in enhancing TNF- $\alpha$  when combined with lenalidomide. Therefore, interstitial lung disease may develop not only due to enhanced secretion of TNF- $\alpha$  by elotuzumab but also because of the synergistic effect of combining elotuzumab with lenalidomide.

## Conclusions

The administration of elotuzumab can cause interstitial pneumonitis. Presentation is quite similar to an infection; differentiation upon radiology is not conclusive and need full infectious workup. Elotuzumab induced interstitial pneumonitis ought to be in the differential diagnosis in patients on elotuzumab combination therapy and presenting with respiratory distress and hypoxic respiratory failure. Treatment with high dose steroids can be useful and should be started early after ruling out an infection.

## Notes

The content published in Cureus is the result of clinical experience and/or research by independent individuals or organizations. Cureus is not responsible for the scientific accuracy or reliability of data or conclusions published herein. All content published within Cureus is intended only for educational, research and reference purposes. Additionally, articles published within Cureus should not be deemed a suitable substitute for the advice of a qualified health care professional. Do not disregard or avoid professional medical advice due to content published within Cureus.

The authors have declared that no competing interests exist.

## Human Ethics

Consent was obtained by all participants in this study

## Article information

Cureus. 2020 Oct; 12(10): e11162.

Published online 2020 Oct 25. doi: [10.7759/cureus.11162](https://doi.org/10.7759/cureus.11162)

PMCID: [PMC7686952](https://pubmed.ncbi.nlm.nih.gov/PMC7686952/)

PMID: [33251069](https://pubmed.ncbi.nlm.nih.gov/33251069/)

Monitoring Editor: Alexander Muacevic and John R Adler

Ghulam Ghous,<sup>1</sup> Hafiz Muhammad Hassan Shoukat,<sup>2</sup> Zahid Ijaz Tarar,<sup>1</sup> and Muhammad Usman Zafar<sup>3</sup>

<sup>1</sup> Internal Medicine, University of Missouri, Columbia, USA

<sup>2</sup> Internal Medicine, Premier Health/Wright State University, Dayton, USA

<sup>3</sup> Hospital Medicine, Lehigh Valley Health Network, Allentown, USA

 Corresponding author.

Ghulam Ghous [ghulam.ghous77@yahoo.com](mailto:ghulam.ghous77@yahoo.com)

Accepted 2020 Oct 25.

Copyright © 2020, Ghous et al.

This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Articles from Cureus are provided here courtesy of **Cureus Inc.**

## References

1. Elotuzumab: a review in relapsed and/or refractory multiple myeloma. Lamb YN. *Drugs*. 2018;78:1481–1488. [[PubMed](#)] [[Google Scholar](#)]
2. Anti-CD38 and anti-SLAMF7: the future of myeloma immunotherapy. Zamagni E, Tacchetti P, Pantani L, Cavo M. *Expert Rev Hematol*. 2018;5:423–435. [[PubMed](#)] [[Google Scholar](#)]
3. Update on elotuzumab for the treatment of relapsed/refractory multiple myeloma: patients' selection and perspective. Trudel S, Moreau P, Touzeau C. *Oncotargets Ther*. 2019;12:5813–5822. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
4. Monoclonal antibodies in relapsed/refractory myeloma: updated evidence from clinical trials, real-life studies, and meta-analyses. Musto P, La Rocca F. *Expert Rev Hematol*. 2020;13:331–349. [[PubMed](#)] [[Google Scholar](#)]
5. Elotuzumab therapy for relapsed or refractory multiple myeloma. Lonial S, Dimopoulos M, Palumbo A, et al. *N Engl J Med*. 2015;373:621–631. [[PubMed](#)] [[Google Scholar](#)]
6. Elotuzumab plus pomalidomide and dexamethasone for multiple myeloma. Dimopoulos MA, Dytfield D, Grosicki S, et al. *N Engl J Med*. 2018;11:1811–1822. [[PubMed](#)] [[Google Scholar](#)]
7. A phase 1, multicenter, open-label, dose escalation study of elotuzumab in patients with advanced multiple myeloma. Zonder JA, Mohrbacher AF, Singhal S, et al. *Blood*. 2012;120:552–559. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
8. Elotuzumab-induced interstitial lung disease: the first case report. Tanaka T, Saito Y, Kokuho N, et al. *Jpn J Clin Oncol*. 2018;48:491–494. [[PubMed](#)] [[Google Scholar](#)]
9. Lenalidomide-associated pneumonitis in patients with plasma cell dyscrasias. Zagouri F, Roussou M, Kastiris E, et al. *Am J Hematol*. 2011;86:882–884. [[PubMed](#)] [[Google Scholar](#)]
10. Drug-induced interstitial pneumonitis due to low-dose lenalidomide. Kunimasa K, Ueda T, Arita M, Maeda T, Hotta M, Ishida T. *Intern Med*. 2012;51:1081–1085. [[PubMed](#)] [[Google Scholar](#)]
11. Lenalidomide-induced eosinophilic pneumonia. Toma A, Rapoport AP, Burke A, Sachdeva A. *Respirol Case Rep*. 2018;48:491–494. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
12. Pomalidomide-induced pulmonary toxicity in multiple myeloma. Modi D, Mamdani H, Vettese T. *Am J Med Sci*. 2015;350:241–242. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
13. Acute lung toxicity related to pomalidomide. Geyer HL, Viggiano RW, Lacy MQ, et al. *Chest*. 2011;140:529–533. [[PubMed](#)] [[Google Scholar](#)]
14. Pulmonary toxicity associated with pomalidomide. Gajic S, Callahan EC, Brown JK, Elicker B, Ley B, Blanc PD. *Int J Tuberc Lung Dis*. 11:1383–1386. [[PubMed](#)] [[Google Scholar](#)]
15. CS1, a potential new therapeutic antibody target for the treatment of multiple myeloma. Hsi ED, Steinle R, Balasa B, et al. *Clin Cancer Res*. 2008;14:2775–2784. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
16. Tumor necrosis factor alpha release is a major biological event associated with rituximab treatment. Bienvenu J, Chvetzoff R, Salles G, et al. *Hematol J*. 2001;2:378–384. [[PubMed](#)] [[Google Scholar](#)]
17. Elotuzumab enhances natural killer cell activation and myeloma cell killing through interleukin-2 and TNF- $\alpha$  pathways. Balasa B, Yun R, Belmar NA, et al. *Cancer Immunol Immunother*. 2015;64:61–73. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]