

Glucose Gel Treatment for Neonatal Hypoglycemia

Juliette Stilley ASN, RN
Lehigh Valley Health Network, Juliette.Stilley@lvhn.org

Ericka E. Clayton BSN, RN
Lehigh Valley Health Network, Ericka_E.Clayton@lvhn.org

Iyanuoluwa Adegbite ASN, RN
Lehigh Valley Health Network, Iyanuoluwa.Adegbite@lvhn.org

Tori Ackley BSN, RN
Lehigh Valley Health Network, Tori.Ackley@lvhn.org

Follow this and additional works at: <https://scholarlyworks.lvhn.org/patient-care-services-nursing>

Let us know how access to this document benefits you

Published In/Presented At

Stilley, J. Clayton, E. Adegbite, I. Ackley, T. (2019, March 28). *Glucose Gel Treatment for Neonatal Hypoglycemia*. Poster presented at: LVHN Vizient/AACN Nurse Residency Program Graduation, Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Glucose Gel Treatment for Neonatal Hypoglycemia

Juliette Stilley ASN, RN, Ericka Clayton BSN, RN, Iyanulouwa Adegbite ASN, RN, Tori Ackley BSN, RN

Lehigh Valley Health Network, Allentown, Pennsylvania

Background/Triggers

- **Transient Neonatal Hypoglycemia:**
 - Hypoglycemia is one of the most commonly encountered problem in infants within the first 48 hours of life
 - At risk infants include: being born late preterm, LGA, SGA, IUGR, and IDM
- **Healthy People 2020 Objectives:**
 - MICH 21.4 and 21.5: increase proportion of infants breastfed exclusively through 3 months/6 months
 - MICH 23: Reduce the proportion of breastfed newborns who receive formula supplementation within the first 2 days of life

Maternal, Infant, and Child Health (MICH).

PICO Question

P: Infant with hypoglycemia
I: Glucose Gel
C: Formula
O: Achieve therapeutic blood sugar levels while still maintaining exclusive breastfeeding, as well as cost effectiveness.

In neonates, does glucose gel compared to formula maintain therapeutic blood sugar levels to promote exclusive breastfeeding?

Evidence


- Reduction of NICU admissions by 73% (Bennett, Fagan, Chaharbakshi, Zamfirova & Flicker, 2016)
- Reduced recurrence rate of hypoglycemia, increased exclusive breastfeeding rates at discharge, and decrease need for admission to the NICU to treat hypoglycemia (Rozance & Hay, 2016)
- Reduce neonatal ICU admissions, reduce hospital length of stay and cost, support the mother/infant dynamic through reduced separation, support exclusive breastfeeding, and improve parental satisfaction (Newman & Bunch, 2017)
- **Gel Cost**
 - 15g tube= 15 doses for a 5kg baby
 - Each tube costs \$2.90
 - Each dose of gel: \$0.19 for 5kg baby

Implementation

- Met with Joyce Najarian, Inpatient DM
 - Informed about project, received no additional information
- Met with Pharmacy
 - Reviewed cost, brand, dosage, and implementation of gel
 - Agreed on pharmacy pre-filling syringes and stocking in pyxis on units
- Mother-Infant Care meeting
 - Presented research to providers
 - Interest shown in future clinical trial
- Proposed pathway



Neonate Birth weight Kg	Dose of 40% Glucose Gel G	Amount mL
2.0	0.4	1
2.5	0.5	1.25
3.0	0.6	1.5
3.5	0.7	1.75
4	0.8	2
4.5	0.9	2.25
5	1	2.5



Lehigh Valley Health Network. (1, May 2017).

Next Steps

- Make TLC to educate staff
- Implement EPIC workspace
- Carry out clinical trial

REFERENCES

1. Center for Disease Control (CDC) (2016). Breastfeeding Report Card: Progressing Toward National Breastfeeding Goals
2. Glasgow, M. J., Harding, J. E., & Edlin, R. (2018). Cost Analysis of Treating Neonatal Hypoglycemia with Dextrose Gel. *Journal Of Pediatrics*, 198151- 155.e1. doi:10.1016/j.jpeds.2018.02.036
3. Lehigh Valley Health Network. (1, May 2017). Assessment of all newborns for prevention & management of hypoglycemia. Retrieved from <https://lvhn.policytech.com/dotNet/documents/?docid=9903>
4. Healthy People 2020. (n.d.). Maternal, Infant, and Child Health (MICH). Retrieved from <https://www.healthypeople.gov/2020/topics-objectives/topic/maternal-infant-and-child-health/objectives>
5. Bennett, C., Fagan, E., Chaharbakshi, E., Zamfirova, I., & Flicker, J. (2016). Implementing a protocol using glucose gel to treat neonatal hypoglycemia. *Nursing for women's health*, 20(1), 64-74.
6. Rozance, P., Hay, H. (2016). New approaches to management of neonatal hypoglycemia. *Maternal Health, Neonatology, and Perinatology* <https://mhnpjournal.biomedcentral.com/articles/10.1186/s40748-016-0031-1>
7. Newman, K., Bunch, M. (2017). Glucose gel as a treatment strategy for transient neonatal Hypoglycemia. *Advances in Neonatal Care*; 17(6) 470-477. <https://www.ncbi.nlm.nih.gov/pubmed/28857766>

© 2014 Lehigh Valley Health Network