

Congenital Nasal Pyriform Aperture Stenosis

Priya P. Dmello BSN, RNC-NIC
Lehigh Valley Health Network, Priya_P.Dmello@lvhn.org

Jane M. Nemeth MSN, RNC-NIC
Lehigh Valley Health Network, Jane_M.Nemeth@lvhn.org

Follow this and additional works at: <https://scholarlyworks.lvhn.org/patient-care-services-nursing>

Let us know how access to this document benefits you

Published In/Presented At

Dmello, P. P., Nemeth, J. M. (2019, Sept 11-14). *Congenital Nasal Pyriform Aperture Stenosis*. Poster Presented at: National Neonatal/Mother Baby/Advanced Practice Nurses Conference, Orlando, FL.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Congenital Nasal Pyriform Aperture Stenosis

Priya Dmello BSN, RNC-NIC and Jane Nemeth, MSN, RNC-NIC
Neonatal Intensive Care Unit, Lehigh Valley Reilly Children's Hospital, Allentown, Pa.


This offering details the presentation, differential diagnosis and evidence-based interprofessional management of a neonate afflicted with Congenital Nasal Pyriform Aperture Stenosis (CNPAS) illustrated by a case study in a level IV neonatal intensive care unit (NICU).

CASE PRESENTATION

- Term female infant delivered via spontaneous vaginal delivery
 - Transferred to mother-baby unit
- Nasal congestion and increased work of breathing (WOB) unrelieved by suctioning at 2–3 hours of life (HOL); transferred to level II NICU
- Difficulty passing NGT on left nares
- Clinical presentation – Noisy breathing, respiratory distress, worsened with feeding/ improved with crying
 - Neonates are obligate nose breathers
 - In CNPAS – overgrowth of anterior bony opening of nose in facial skeleton causes airway obstruction
- ABG and CXR within normal limits (wnl)
- Required 3L HFNC; weaned to oxyhood with cool mist
- Day of life (DOL) 2 started on Neosynephrine nasal drops q12 hours
- DOL 3 – able to pass NGT bilaterally; however, continued increased WOB transferred to level IV NICU for higher level of care and ENT evaluation


DIAGNOSTIC FINDINGS

- ENT unable to pass 3.5mm flexible scope past anterior right nares
- CT scan face- confirmed CNPAS
- Discovered single central maxillary incisor (may be associated with pituitary abnormalities)
- Brain MRI – normal pituitary, no holoprosencephaly
- Pediatric Endocrine consult- often high association with growth hormone deficiency
 - Glucoses wnl x 24hrs
 - IGF-1 wnl
- ECHO in light of other midline anomalies – small VSD
 - Microarray wnl



UNCOVERING THE DIAGNOSIS

- Inability/difficulty passing NGT or bedside scope
- CT scan:
 - Small nasal opening



Left - normal infant
Right - infant with CNPAS- narrowed opening of nasal aperture



Axial view showing single central maxillary incisor
Yang, Sara et al. "Congenital nasal pyriform aperture stenosis in association with solitary median maxillary central incisor: Unique radiologic features." *Radiology Case Reports* vol. 11,3 178-81. 1 Aug. 2016. doi:10.1016/j.radcr.2016.06.004

STANDARD TREATMENT

- Recommend endocrine consult and baseline endocrine labs for pituitary evaluation
 - IGF-1, cortisol, FT4 and TSH
- Management – ENT consult
- Conservative therapy – suctioning, positioning, parenteral/ enteral support
 - More severe forms usually require surgical correction (Dilation and/or drilling, nasal stenting)
- Decongestants – often sufficient for patients with mild CNPAS
 - Nasal saline drops (gold standard)
 - Dexamethasone 0.1% ophthalmic solution intranasally (2nd line)
 - Phenylephrine 0.125% nasal drops (3rd line)

HOSPITAL COURSE

- Started on intranasal dexamethasone drops
- Airway positional requiring side-lying placement; improved by DOL 4, able to maintain O2 saturation levels supine
- D/c to home on DOL 5 to complete 10-day dexamethasone nasal drops course
- Continued to have nasal congestion and intermittent increased WOB
- 3 weeks of life (WOL) – underwent pyriform aperture stenosis drill out with dilation and stent placement
- Nasal stents removed on post-op day (POD) #2 and D/c to home with 10-day course of dexamethasone nasal drops
- 7 WOL – underwent nasal endoscopy with dilation and stent placement
- Stents removed on POD #2 – D/c home with dexamethasone drops and ofloxacin drops



Rajan, R., and Tunkel, D. (2018). Choanal atresia and other neonatal nasal anomalies. *Clinics in Perinatology*, Volume 45(4), 751 – 767.

NURSING PEARLS

- Watch for hypoglycemia, hypotension, conjugated/unconjugated hyperbilirubinemia
- Interprofessional collaboration is essential (speech, PT, OT, medicine-endocrine/ENT/Neonatology)
- Family support is crucial- need assistance with feeding, positioning, coordination of care and psychosocial support

REFERENCES

1. Rajan, R., and Tunkel, D. (2018). Choanal atresia and other neonatal nasal anomalies. *Clinics in Perinatology*, Volume 45(4), 751 – 767.
2. Sesenna, E., Leporati, M., Brevi, B., Oretti, G., and Ferri, A. (2012). Congenital nasal pyriform aperture stenosis: Diagnosis and management. *Italian Journal of Pediatrics*, 38(28). doi:10.1186/1824-7268-38-28
3. Yang, Sara et al. (2016). Congenital nasal pyriform aperture stenosis in association with solitary median maxillary central incisor: Unique radiologic features. *Radiology Case Reports* vol. 11(3) 178-81.1 doi:10.1016/j.radcr.2016.06.004