

Clinical Indications for Peripheral IV Catheter Replacements

Liudmila Fedor RN BSN
Lehigh Valley Health Network, Liudmila.Fedor@lvhn.org

Charlene Langley RN

Callie Waldron RN BSN
Lehigh Valley Health Network, Callie.Waldron@lvhn.org

Follow this and additional works at: <https://scholarlyworks.lvhn.org/patient-care-services-nursing>

Let us know how access to this document benefits you

Published In/Presented At

Fedor, L. Langley, C. Waldron, C. (2019, November 7). *Clinical Indications for Peripheral IV Catheter Replacements*. Poster Presented at: LVHN Vizient/AACN Nurse Residency Program Graduation, Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Clinical Indications for Peripheral IV Catheter Replacements

6B Nurse Residents: Liudmila Fedor RN BSN, Charlene Langley RN, Callie Waldron RN BSN

Lehigh Valley Health Network, Allentown, Pennsylvania

BACKGROUND

- Current practice at LVHN is to replace IV catheters after 72 hours regardless of if the site is symptomatic or not. Current practice also causes them unnecessary pain while having a new IV placed when not clinically indicated.
- Renal patients are usually harder to obtain IV access in, changing sites when clinically indicated increases patient satisfaction and decreases number of sticks the patient has to have.

PICO


- **P – Adult Medical-Surgical Patients**
- **I – Clinically indicated replacement of IV catheters**
- **C – Current Practice – 72hours**
- **O – Impact on IV associated complications**

EVIDENCE

- Rate of actual IV associated blood stream infections is only 0.5 per 1000 catheter days.
- Many patients are afraid of needles and will refuse IV changes unless really necessary.
- No significant changes found among 755 inpatients when comparing clinically indicated changes versus routine changes.
- In a systematic review of 7 trials with a total of 4895 patients, no evidence was found to support changing catheters every 72-96 hours (Webster, Osborne, Rickard, New, 2015.)

OUTCOMES

- Infection rates are the same when IVs are changed routinely versus when clinically indicated.
- The number of symptomatic IV sites does not increase if the IV is in for longer than 72 hours.
- Less frequent IV changes increases patient satisfaction.
- Less IV changes decreases hospital expenses on IV catheters



IMPLEMENTATION

- Assess site routinely for signs of infection, phlebitis, and infiltration
- Flush IV catheters at least once a shift to ensure patency.
- Change catheters when clinically indicated instead of every 72 hours.
- Survey staff to assess why IV sites are being changed.

NEXT STEPS

- This project was not approved for clinical implementation due to associated risks and disagreement with current network practices.
- Staff survey was not handed out due to the project not being approved by the network.
- Another alternative would be to modify the current LVHN practice guidelines to allow IV catheter replacement every 96 hours compared to the current 72 hours.

REFERENCES

- Brown, D., & Rowland, K. (2013). PURLs: optimal timing for peripheral IV replacement?. *The Journal of family practice*, 62(4), 200–202.
- Fang, C. (2012, September). Peripheral IV in Too Long. Retrieved from <https://psnet.ahrq.gov/webmm/case/278/Peripheral-IV-in-Too-Long>
- Recommendations | BSI | Guidelines Library | Infection Control | CDC. (2015, November 5). Retrieved April 5, 2019, from <https://www.cdc.gov/infectioncontrol/guidelines/bsi/recommendations.html>
- Webster J, Osborne S, Rickard CM, New K. Clinically - indicated replacement versus routine replacement of peripheral venous catheters. *Cochrane Database of Systematic Reviews* 2015, Issue 8. Art. No.: CD007798. DOI: 10.1002/14651858.CD007798.pub4.
- Xu, L., Hu, Y., Huang, X., Fu, J., & Zhang, J. (2017). Clinically indicated replacement versus routine replacement of peripheral venous catheters in adults: A nonblinded, cluster-randomized trial in China. *International Journal Of Nursing Practice*, 23(6). <https://doi.org/10.1111/ijn.12595>