

The CIWA Assessment and Education

Jenna Lukach RN

Lehigh Valley Health Network, jenna.lukach@lvhn.org

Alessandra Pagnotta RN

Lehigh Valley Health Network, Alessandra.Pagnotta@lvhn.org

Haley Ragazinsky RN

Lehigh Valley Health Network, Haley.Ragazinsky@lvhn.org

Follow this and additional works at: <https://scholarlyworks.lvhn.org/patient-care-services-nursing>



Part of the [Nursing Commons](#)

Published In/Presented At

Lukach, J., Pagnotta, A., & Ragazinsky, H. (2020, August). The CIWA Assessment and Education. Poster presented at LVHN Vizient/AACN Nurse Residency Program Graduation, Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

The CIWA Assessment and Education

Jenna Lukach, RN, Alessandra Pagnotta, RN, Haley Ragazinsky, RN

Lehigh Valley Health Network, Allentown, Pennsylvania

BACKGROUND

- The CIWA assessment is a tool used to assess patients withdrawing from alcohol. This assessment results in a dosing-based score from a variety of patient symptoms.
- CIWA scoring is new to LVHN-Schuylkill and it was brought to our attention that nurses had a lack of knowledge related to CIWA.
- Alcohol withdrawal is seen frequently on the psychiatric unit and medical-surgical units

PICO


- **P:** 30 RN's on med-surg, telemetry, and psychiatric units
- **I:** Education packet that provided knowledge of CIWA
- **C:** A brief TLC was provided on CIWA when it was first introduced at LVH-Schuylkill
- **O:** Scores on post-test after further education was provided

EVIDENCE

- More than 16 million people in the U.S. are affected by alcoholism (Eloma et al., 2018)
- Symptoms start 4-72 hours after cessation of drinking. (Eloma et al., 2018)
- CIWA protocol allows for a shorter period of detoxification, lower doses of benzodiazepines, less sedation effects, and a decreased chance of respiratory depression. (Eloma et al., 2018)

OUTCOMES

- The pre and post tests were identical, 10 question tests. The following scores were:



IMPLEMENTATION

- Knowledge reinforcement in the form of educational packets were given after the pre-test.
- Education included signs and symptoms of alcohol withdrawal, background of information about CIWA, and assessment/treatment of alcohol withdrawal.

NEXT STEPS

- Continue educating on symptoms of alcohol withdrawal and treatment and allow nurses to teach back the education provided.
- Use of CIWA is vital to ensure the patients who require this type of treatment are getting the highest quality care possible.

REFERENCES:

- Eloma, A., Tucciarone, J., Hayes, E., & Bronson, B. (2018). Evaluation of the appropriate use of a CIWA-Ar alcohol withdrawal protocol in the general hospital setting. *American Journal of Drug and Alcohol Abuse*, 44(4), 418-425. Retrieved from EBSCOhost database. (Accession No. 129527864)
- Higgins, J., Bugajski, A., Church, D., Oylar, D., Parli, S., Halcomb, P., . . . Bernard, A. (2019). A psychometric analysis of CIWA-Ar in acutely ill and injured hospitalized patients. *Journal of Trauma Nursing*, 26(1), 41-49. <https://doi.org/10.1097/JTN.0000000000000414>
- Jesse, S., Bråthen, G., Ferrara, M., Keindl, M., Ben-Menachem, E., Tanasescu, R., . . . Ludolph, A. C. (2016, September 1). Alcohol withdrawal syndrome: mechanisms, manifestations, and management. Retrieved from <https://onlinelibrary.wiley.com/doi/epdf/10.1111/ane.12671>
- Naveen Kumar, C., Andrade, C., & Murthy, P. (2009). A randomized, double-blind comparison of lorazepam and chlordiazepoxide in patients with uncomplicated alcohol withdrawal. *Journal of Studies on Alcohol and Drugs*, 70(3). <https://doi.org/10.15288/jsad.2009.70.467>