Lehigh Valley Health Network

LVHN Scholarly Works

Department of Medicine

AXIOS Lumen Apposing Metal Stent in the Treatment of Peripancreatic Fluid Collections, 4 Year Retrospective Observation at a Large Care Center

Michal Kloska MD Lehigh Valley Health Network, Michal.Kloska@lvhn.org

Valery Hrad MD Lehigh Valley Health Network, Valery. Hrad@lvhn.org

Abdul Aleem MD Lehigh Valley Health Network, Abdul.Aleem@lvhn.org

Soorya N. Aggarwal DO

Shashin Shah MD Lehigh Valley Health Network, Shashin.Shah@lvhn.org

See next page for additional authors

Follow this and additional works at: https://scholarlyworks.lvhn.org/medicine



Part of the Gastroenterology Commons

Let us know how access to this document benefits you

Published In/Presented At

Kloska, M., Hrad, V., Aleem, A., Aggarwal, S., Shah, S., Zator, Z., & Shah, H. (2020, October). AXIOS Lumen Apposing Metal Stent in the Treatment of Peripancreatic Fluid Collections, 4 Year Retrospective Observation at a Large Care Center. Poster presented at: ACG Annual Scientific Meeting, Virtual.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

authors	
Aichal Kloska MD, Valery Hra ator MD, and Hiral N. Shah I	ad MD, Abdul Aleem MD, Soorya N. Aggarwal DO, Shashin Shah MD, Zacha MD

AXIOS Lumen Apposing Metal Stent in the Treatment of Peripancreatic Fluid Collections, 4 Year Retrospective Observation at a Large Care Center

Michal Kloska, MD, PhD,¹ Valery Hrad, MD,² Abdul Aleem, MD,² Soorya Aggarwal, DO,³ Shashin Shah, MD,² Zachary Zator, MD,² Hiral N. Shah, MD¹

¹Department of Medicine, ²Department of Gastroenterology, Lehigh Valley Health Network, Allentown, Pa., ³Department of Gastroenterology, MedStar Georgetown University Hospital, Washington, DC

Introduction

- Peripancreatic fluid collections (PFCs) are one of the most common complications of acute pancreatitis.
- The majority of PFCs resolve spontaneously, the rest form mature cysts filled with either necrotic debris or with fluid, classified as walled off pancreonecrosis (WOPN) and pancreatic pseudocysts (PP) respectively.
- Electrocautery enhanced AXIOS™ lumenapposing metal stents (EEAL) has primed EUS guided drainage as a treatment of choice for PFCs.

Methods

- IRB approved retrospective chart review of EEAL performed at a large quaternary care center.
- Procedural data including technical and clinical success rate were collected and analyzed.
- Clinical success rate was measured by symptoms resolution at first outpatient office visit and by cyst resolution (complete, incomplete (>50%) or partial (<50%).

Results


- 38 patients underwent 51 EEAL procedures for the treatment of PFCs.
- 92.2% of the cysts were at least 6cm in diameter.
- 11 patients required multiple EEAL placement, of whom 7 patients had multiple cysts requiring separate treatments, 2 patients had reoccurrence of the cysts requiring drainage, 1 patient required 2 EEAL placed for one large septated cyst and 1 patient required 2 attempts to place the stent.

- Technical success rate was 96.1%, 2 times EEAL angulated during placement (stent dislodgment to the stomach, stent placement to the fluid surrounding the PFCs)
- Only after 5 interventions (9.8%) patients didn't experience any clinical improvement postprocedurally of which 2 required surgical intervention, 2 had cyst reoccurrence and 1 had stent placement in the pericystic area.
- The average hospital stay after EEAL placement was 7.89 days with a mean of 3.5 days.

Conclusion

Our data demonstrates safety and efficacy of EEAL use in the treatment of PFCs with a high technical success rate of 96.1%. Over 90% of patients experienced partial or complete resolution of their symptoms and 83.4% had at least 50% of cyst size reduction was noted. Larger studies in this area is needed to prove EEAL's superiority over more traditional methods of treating PFCs.

Mean age	54.5 (22-75) yo
Gender	Male 71% Female 29%
Infected necrosis	37.3%
PFC reoccurrence	12.5%
Surgical intervention	3.9%



6-8.9cm

>9cm

